

# USTEKINUMAB

(Including Stelara and biosimilars: Wezlana, Yesintek, Selarsdi, Pyzchiva, Steqeyma)

**FLEXCARE**  
INFUSION CENTERS

## PATIENT DEMOGRAPHICS

Patient Name:	Patient's Phone Number:
Date of Birth:	Address:
Allergies: See List <input type="checkbox"/> NKDA <input type="checkbox"/>	City, State, Zip:
Weight: _____ lbs or _____ kg	Patient's Email:

## REQUIRED DOCUMENTATION

• Insurance Card • H&P • Patient Demographics • Most Recent Labs • Med List • Tried/Failed Therapies • Neg TB Results

## PRIMARY DIAGNOSIS

- |   |   |
|---|---|
| <input type="checkbox"/> K50.00 Crohn's disease of small intestine without complications (CD) | <input type="checkbox"/> K51.90 Ulcerative colitis, unspecified, w/o complications (UC) |
| <input type="checkbox"/> K50.019 Crohn's disease of small intestine with unsp comp (CD)       | <input type="checkbox"/> L40.5 Psoriatic Arthritis (PsA)                                |
| <input type="checkbox"/> K50.10 Crohn's disease of large intestine without complications (CD) | <input type="checkbox"/> L40.9 Plaque Psoriasis (Ps)                                    |
| <input type="checkbox"/> K50.90 Crohn's disease, unspecified, without complications (CD)      | <input type="checkbox"/> Other: _____   |
| <input type="checkbox"/> K51.00 Ulcerative (chronic) pancolitis without complications (UC)    | _____   |

## LAB ORDERS: PLEASE INCLUDE FREQUENCY

Please list any labs to be drawn by the infusion clinic: \_\_\_\_\_

## PRE-MEDICATIONS

\*Per infusion clinic protocol: No recommended standard pre-meds for Stelara

☐ Provider Prescribed: \_\_\_\_\_

## PRIMARY MEDICATION ORDER

\*Biosimilar (Wezlana, Yesintek, Selarsdi, Pyzchiva, Steqeyma) may be used according to payer guidelines

\*To prohibit auto-substitution, please indicate specific brand required: \_\_\_\_\_

### Ulcerative Colitis (UC) – or – Crohn's Disease (CD)

Induction Doses (to be administered in infusion clinic):

- ☐ Weight <55kg: Stelara 260mg IV once  
☐ Weight 55kg: Stelara 390mg IV once  
☐ Weight >85kg: Stelara 520mg IV once

Maintenance Doses:

- ☐ Infusion clinic will coordinate with SP for self-administration or administration in-clinic as payor dictates: Stelara 90mg subQ every 8 weeks after induction dose.  
☐ Provider's office will coordinate initial maintenance dose from SP.

### Plaque Psoriasis (Ps) – or – Psoriatic Arthritis (PsA)

- ☐ Weight ≤ 100kg: Stelara 45mg subQ at Weeks 0, 4, and every 12 weeks thereafter  
☐ Weight > 100kg: Stelara 90mg subQ at Weeks 0, 4, and every 12 weeks thereafter

\*Infusion clinic will coordinate initial dose from Specialty Pharmacy

☐ Other: \_\_\_\_\_

First Dose: ☐ Y ☐ N ☒ Refill x12 months unless otherwise noted: \_\_\_\_\_

## LINE USE/CARE ORDERS

☒ Start PIV/ACCESS CVC ☒ Flush device per FlexCare Infusion Centers' protocol (See flexcareinfusion.com for detailed policy)

☐ Other Flush Orders: Please fax other line care orders if checking this box

## ADVERSE REACTION & ANAPHYLAXIS ORDERS

☒ Administer acute infusion reaction and anaphylaxis medications per FlexCare Infusion Centers' protocol (See flexcareinfusion.com for detailed policy) ☐ Other: Please fax other reaction orders if checking this box

## PROVIDER INFORMATION: PLEASE CHECK PREFERRED FORM OF COMMUNICATION

Provider Name:	Office Contact:
Address:	Phone:
City, State, Zip:	<input type="checkbox"/> Fax:
NPI AND License:	<input type="checkbox"/> Email:

Provider Signature

Date

FAX: (888) 219-8102 | EMAIL: [orders@flexcareinfusion.com](mailto:orders@flexcareinfusion.com) | VISIT: [flexcareinfusion.com/referrals](http://flexcareinfusion.com/referrals)